

A SHORT PROOF OF HRS-TILTING

XIAO-WU CHEN

ABSTRACT. We give a short proof to the following tilting theorem by Happel, Reiten and Smalø via an explicit construction: given two abelian categories $\mathcal{A}$ and $\mathcal{B}$ such that $\mathcal{B}$ is tilted from $\mathcal{A}$, then $\mathcal{A}$ and $\mathcal{B}$ are derived equivalent.

1. INTRODUCTION

Let $\mathcal{A}$ be an abelian category. Recall that a *torsion pair* on $\mathcal{A}$ means a pair $(\mathcal{T}, \mathcal{F})$ of full subcategories satisfying

- (T1). $\text{Hom}_{\mathcal{A}}(T, F) = 0$ for all $T \in \mathcal{T}$ and $F \in \mathcal{F}$; both subcategories $\mathcal{T}$ and $\mathcal{F}$ are closed under direct summands;
- (T2). for each object $X \in \mathcal{A}$, there is a short exact sequence $0 \longrightarrow T \longrightarrow X \longrightarrow F \longrightarrow 0$ for some $T \in \mathcal{T}$ and $F \in \mathcal{F}$.

In a torsion pair $(\mathcal{T}, \mathcal{F})$, it follows that the subcategory $\mathcal{T}$ is closed under extensions and factor objects; $\mathcal{F}$ is closed under extensions and sub objects. The torsion pair $(\mathcal{T}, \mathcal{F})$ is called a *tilting torsion pair*, provided that each object in $\mathcal{A}$ embeds into an object in $\mathcal{T}$. Dually the torsion pair $(\mathcal{T}, \mathcal{F})$ is called a *cotilting torsion pair*, provided that each object in $\mathcal{A}$ is a factor object of an object in $\mathcal{F}$ ([3, Chapter I, section 3]).

Denote a complex in $\mathcal{A}$ by $X^\bullet = (X^n, d_X^n)_{n \in \mathbb{Z}}$ where $d_X^n : X^n \longrightarrow X^{n+1}$ is the differential satisfying $d_X^{n+1} \circ d_X^n = 0$; its shift $X^\bullet[1]$ is a complex given by $(X^\bullet[1])^n = X^{n+1}$ and $d_{X[1]}^n = -d_X^{n+1}$. Denote by $D(\mathcal{A})$ the (unbounded) *derived category* of $\mathcal{A}$, $D^+(\mathcal{A})$, $D^-(\mathcal{A})$ and $D^b(\mathcal{A})$ the full subcategory consisting of bounded-below, bounded-above and bounded complexes, respectively ([6, 4]). We will always identify the abelian category $\mathcal{A}$ as the full subcategory of $D(\mathcal{A})$ consisting of stalk complexes concentrated at degree zero ([4, p.40, Proposition 4.3]). Let $(\mathcal{T}, \mathcal{F})$ be a torsion pair on $\mathcal{A}$. Following [3, Chapter I, section 2], set $\mathcal{B}$ to be the full subcategory of $D(\mathcal{A})$ consisting of complexes $X^\bullet$ satisfying $H^0(X^\bullet) \in \mathcal{T}$, $H^{-1}(X^\bullet) \in \mathcal{F}$ and $H^i(X^\bullet) = 0$ for $i \neq 0, 1$. Note that $\mathcal{T} \subseteq \mathcal{B}$ and $\mathcal{F}[1] \subseteq \mathcal{B}$. By [3, Chapter I, Proposition 2.1] the category $\mathcal{B}$ is the heart of certain t -structure on $D(\mathcal{A})$ and thus by [1] it is an abelian category (also see [2]); moreover the pair $(\mathcal{F}[1], \mathcal{T})$ is a torsion pair on $\mathcal{B}$.

One might expect that the resulting new abelian category $\mathcal{B}$ is derived equivalent to $\mathcal{A}$. But this is in general false (by the example in [3, p.16]). However Happel, Reiten and Smalø show the following remarkable result ([3, Chapter I, Theorem 3.3]).

Theorem (Happel-Reiten-Smalø) *Let $(\mathcal{T}, \mathcal{F})$ be a tilting torsion pair on $\mathcal{A}$. Then we have a natural equivalence of triangulated categories $D(\mathcal{B}) \simeq D(\mathcal{A})$ which is compatible with the inclusion of $\mathcal{B}$ into $D(\mathcal{A})$. Similar results hold for $D^*(-)$ with $*$ in $\{+, -, b\}$.*

Key words and phrases. torsion pair, tilting, derived equivalence.

This project was supported by Alexander von Humboldt Stiftung, and was also partially supported by China Postdoctoral Science Foundation No. 20070420125 and No. 200801230. The author also gratefully acknowledges the support of K. C. Wong Education Foundation, Hong Kong.

E-mail: xwchen@mail.ustc.edu.cn.

In the case of Theorem the category $\mathcal{B}$ is said to be *tilted* from $\mathcal{A}$. Note that the original theorem only claims the equivalence between the bounded derived categories and requires the existence of enough projective or injective objects. The quoted version is improved by Noohi ([5, Theorem 7.6]). We will give a short proof of the theorem via an explicit construction of the equivalence functor.

2. THE PROOF OF THEOREM

Throughout $(\mathcal{T}, \mathcal{F})$ is a torsion pair on $\mathcal{A}$ and $\mathcal{B}$ is the resulting abelian category. We start with an easy observation.

Lemma 2.1. *Consider an complex $T^\bullet = (T^n, d_T^n)_{n \in \mathbb{Z}}$ with terms in $\mathcal{T}$. Then it is exact in $\mathcal{A}$ if and only if it is exact in $\mathcal{B}$.*

Proof. Assume that $T^\bullet$ is exact in $\mathcal{A}$. Since $\mathcal{T}$ is closed under factor objects in $\mathcal{A}$, the complex $T^\bullet$ splits into short exact sequences $\xi^n : 0 \rightarrow T^n \xrightarrow{i^n} T^n \xrightarrow{p^n} T^{n+1} \rightarrow 0$ with $T^n \in \mathcal{T}$ and $d_T^n = i^{n+1} \circ p^n$. Since $\mathcal{B}$ is a heart of certain t -structure on $D(\mathcal{A})$, a sequence $0 \rightarrow B^0 \xrightarrow{f} B^1 \xrightarrow{g} B^2 \rightarrow 0$ in $\mathcal{B}$ is short exact if and only if there is a triangle $B^0 \xrightarrow{f} B^1 \xrightarrow{g} B^2 \rightarrow B^0[1]$ in $D(\mathcal{A})$ ([1] and [2, Chapter IV, §4]). Note that short exact sequences in $\mathcal{A}$ induces triangles in $D(\mathcal{A})$ ([4, p.62, Proposition 6.1]). Hence ξ^n become short exact sequences in $\mathcal{B}$. Thus by splicing them together we show that the complex $T^\bullet$ is exact in $\mathcal{B}$. The “if” part is proved similarly. ■

The following result is needed.

Lemma 2.2. (the “only if” part of [3, Chapter I, Proposition 3.2 (i)]) *Let $(\mathcal{T}, \mathcal{F})$ be a torsion pair on $\mathcal{A}$ and let $\mathcal{B}$ as before. If the torsion pair $(\mathcal{T}, \mathcal{F})$ is tilting, then the resulting torsion pair $(\mathcal{F}[1], \mathcal{T})$ on $\mathcal{B}$ is cotilting.*

Remark 2.3. Note that the converse of the above lemma is also true as stated in [3, Chapter I, Proposition 3.2 (i)]. However it seems to the author that a dual argument of this lemma is not working. Instead, thanks to Theorem and then by combining [3, Chapter I, Proposition 3.4] and the “only if” part of [3, Chapter I, Proposition 3.2 (ii)] one deduces that the converse holds (here one needs the fact that the equivalence in Theorem is compatible with the inclusion $\mathcal{B} \hookrightarrow D(\mathcal{A})$).

Proof of Theorem: Denote by $K(\mathcal{A})$ the homotopy category of complexes in $\mathcal{A}$, $K(\mathcal{T})$ (resp. $K_{\text{ex}}(\mathcal{A})$) its full subcategory consisting of complexes in $\mathcal{T}$ (resp. exact complexes). The inclusion $K(\mathcal{T}) \hookrightarrow K(\mathcal{A})$ induces the following exact functor

$$F : K(\mathcal{T})/K(\mathcal{T}) \cap K_{\text{ex}}(\mathcal{A}) \rightarrow D(\mathcal{A}).$$

Since the torsion pair $(\mathcal{T}, \mathcal{F})$ is tilting and $\mathcal{T}$ is closed under factor objects, we have for each $X \in \mathcal{A}$ a short exact sequence $0 \rightarrow X \rightarrow T^0 \rightarrow T^1 \rightarrow 0$ with $T^i \in \mathcal{T}$. Note further that $\mathcal{T}$ is closed under extensions, we infer that the conditions in [4, p.42, Lemma 4.6 2)] are fulfilled, and thus for each complex $X^\bullet$ in $K(\mathcal{A})$ there is a quasi-isomorphism $X^\bullet \rightarrow T^\bullet$ with $T^\bullet \in K(\mathcal{T})$. This implies that the functor F is dense and by [6, p.283, 4-2 Théorème] it is fully-faithful, that is, the functor F is an equivalence of triangulated categories. By Lemma 2.2 we may apply the dual argument to obtain a natural equivalence

$$G : K(\mathcal{T})/K(\mathcal{T}) \cap K_{\text{ex}}(\mathcal{B}) \rightarrow D(\mathcal{B}).$$

By Lemma 2.1 we have $K(\mathcal{T}) \cap K_{\text{ex}}(\mathcal{A}) = K(\mathcal{T}) \cap K_{\text{ex}}(\mathcal{B})$. Hence $FG^{-1} : D(\mathcal{B}) \rightarrow D(\mathcal{A})$ is the required equivalence, where G^{-1} denotes a quasi-inverse of G .

To see other equivalences, let $*$ $\in \{+, -, b\}$ and let $K^*(-)$ denote the corresponding homotopy categories. Note that in the argument above, for a complex $X^\bullet \in K^*(\mathcal{A})$ we may take a quasi-isomorphism $X^\bullet \rightarrow T^\bullet$ with $T^\bullet \in K^*(\mathcal{T})$ (for the case $*$ = +, just consult

the proof in [4, p.43, 1]); for the case $*$ = $-$, because $\mathcal{T}$ is closed under factor objects one may replace $T^\bullet$ by its good truncations; for the case $*$ = b , consult the proof in [4, p.43, 1]) and note that since $\mathcal{T}$ is closed under factor objects, the argument therein is done within finitely many steps, consequently the obtained complex $T^\bullet$ is bounded). Thus we construct the equivalences F^* and G^* as above. This proves the corresponding equivalences between the derived categories $D^*(-)$.

Finally we will show that the obtained equivalence FG^{-1} is compatible with the inclusion $\mathcal{B} \hookrightarrow D(\mathcal{A})$. This is subtle. Given an object $B \in \mathcal{B}$, since the torsion pair $(\mathcal{F}[1], \mathcal{T})$ is cotilting, we have a short exact sequence in $\mathcal{B}$, $\eta : 0 \rightarrow T^{-1} \xrightarrow{d} T^0 \xrightarrow{g} B \rightarrow 0$ with $T^i \in \mathcal{T}$, in other words, a triangle $\xi : T^{-1} \xrightarrow{d} T^0 \xrightarrow{g} B \rightarrow T^{-1}[1]$ in $D(\mathcal{A})$. Then by construction $FG^{-1}(B)$ is isomorphic to the complex $T^\bullet = \cdots \rightarrow 0 \rightarrow T^{-1} \xrightarrow{d} T^0 \rightarrow 0 \rightarrow \cdots$. Note that the complex $T^\bullet$ is the mapping cone of d and thus from the triangle ξ we obtain $T^\bullet$ is isomorphic to B ([4, p.23, Proposition 1.1 c)), in particular $T^\bullet \in \mathcal{B}$. Note the following natural triangle $T^{-1} \xrightarrow{d} T^0 \rightarrow T^\bullet \rightarrow T^{-1}[1]$ and thus a short exact sequence $\gamma : 0 \rightarrow T^{-1} \xrightarrow{d} T^0 \rightarrow T^\bullet \rightarrow 0$ in $\mathcal{B}$. Comparing the short exact sequences η and γ we obtain a unique isomorphism $\theta_B : B \simeq T^\bullet$ in $\mathcal{B}$. We claim that θ is natural in B and then we obtain a natural isomorphism between the inclusion functor $\mathcal{B} \hookrightarrow D(\mathcal{A})$ and the composite $\mathcal{B} \hookrightarrow D(\mathcal{B}) \xrightarrow{FG^{-1}} D(\mathcal{A})$ (here we identify $T^\bullet$ with $FG^{-1}(B)$).

In fact, given a morphism $f : B \rightarrow B'$ in $\mathcal{B}$, choose an exact sequence $\eta' : 0 \rightarrow T'^{-1} \xrightarrow{d'} T'^0 \xrightarrow{g'} B' \rightarrow 0$ with $T'^i \in \mathcal{T}$. Form the complex $T'^\bullet$ and then obtain the short exact sequence $\gamma' : 0 \rightarrow T'^{-1} \xrightarrow{d'} T'^0 \rightarrow T'^\bullet \rightarrow 0$ in $\mathcal{B}$. Identify $G(T^\bullet)$ with B , $G(T'^\bullet)$ with B' . Since the functor G is fully-faithful, we have a chain map $\phi^\bullet : T^\bullet \rightarrow T'^\bullet$ such that $G(\phi^\bullet) = f$. This implies the following commutative exact diagram in $\mathcal{B}$

$$\begin{array}{ccccccc} 0 & \longrightarrow & T^{-1} & \xrightarrow{d} & T^0 & \xrightarrow{g} & \longrightarrow 0 \\ & & \downarrow \phi^{-1} & & \downarrow \phi^0 & & \downarrow f \\ 0 & \longrightarrow & T'^{-1} & \xrightarrow{d'} & T'^0 & \xrightarrow{g'} & B' \longrightarrow 0. \end{array}$$

From this it is direct to see that $\theta_{B'} \circ f = \phi^\bullet \circ \theta_B$ in $\mathcal{B}$ and thus in $D(\mathcal{A})$. This finishes the proof. $\blacksquare$

REFERENCES

- [1] A. BEILINSON, J. BERNSTEIN, P. DELIGNE, Faisceaux pervers, Astérisque **100**, Société Mathématique de France, Paris, 1982.
- [2] S.I. GELFAND, YU I. MANIN, Methods of homological algebra, Springer, 1996.
- [3] D. HAPPEL, I. REITEN, S.O. SMALØ, Tilting in Abelian Categories and Quasitilted Algebras, Memoirs Amer. Math. Soc., vol. **120** no. 575, Providence, Rhode Island, 1996.
- [4] R. HARTSHORNE, Residue and Duality, Lecture Notes in Math. **20**, Springer-Verlag, 1966.
- [5] B. NOOHI, *Explicit HRS-tilting*, Journal of Noncommutative Geometry, to appear.
- [6] J.L. VERDIER, Catégories dérivées, état 0, Springer Lecture Notes **569** (1977), 262-311.

Xiao-Wu Chen, Department of Mathematics, University of Science and Technology of China, Hefei 230026, P. R. China

Current address: Institut fuer Mathematik, Universitaet Paderborn, 33095, Paderborn, Deutschland